

E-Cigarettes Expose NC Youth to the Health Risks of Nicotine

Did you know...

Nicotine is more harmful than you think. **NO AMOUNT IS SAFE FOR YOUTH.**



Among high school students, e-cigarette use is now almost double conventional cigarette use. Nearly all e-cigarettes contain nicotine.^[11]

Nicotine is a chemical commonly found in cigarettes, e-cigarettes and other tobacco products. **Nearly one in seven of North Carolina high school students who has tried e-cigarettes has never smoked a cigarette.**^[2] **A growing body of evidence from multiple countries shows that young people who have never smoked cigarettes — but currently use e-cigarettes — are more likely to smoke cigarettes in the future than are young people who do not use e-cigarettes.**^[12-19] Nicotine is highly addictive and can be toxic.

■ No amount of nicotine is safe for young people.

Nicotine can harm the brain as it continues to develop through young adulthood until age 25.

Animal research has found that even in small doses, nicotine exposure causes long-lasting changes in brain development. The kind of changes that can happen affect learning, memory and addiction to nicotine.^[1-5]

■ Nicotine is harmful to the health of unborn children.

Evidence shows that fetal exposure to nicotine can have negative long-term effects, including sudden infant death syndrome (SIDS), impaired fetal brain and lung development, hearing problems, effects on behaviors and obesity, and deficits in attention and cognition.^[6]

Studies also indicate that fetal nicotine exposure is associated with nicotine dependence in adolescence.^[6-8]

■ Nicotine can be toxic, even deadly.

Eating, drinking or absorbing nicotine can lead to nicotine poisoning; children are especially vulnerable.^[6] Symptoms of nicotine poisoning include nausea, vomiting, seizures and respiratory depression.^[9-10] Nicotine poisoning can be deadly.

For poison emergencies or questions, call the Carolinas Poison Center at 1-800-222-1222.

Learn more about e-cigarettes at www.tobaccopreventionandcontrol.ncdhhs.gov/ecigs.



State of North Carolina • Department of Health and Human Services
Division of Public Health • Tobacco Prevention & Control Branch

www.ncdhhs.gov • NCDHHS is an equal opportunity employer and provider. 04/2018
15,500 copies of this document were printed at a cost of \$1,151.51 or \$0.074291 per copy.

REFERENCES

1. Counotte et al., "Long-Lasting Cognitive Deficits Resulting from Adolescent Nicotine Exposure in Rats."
2. Abreu-Villaca et al., "Short-Term Adolescent Nicotine Exposure Has Immediate and Persistent Effects on Cholinergic Systems: Critical Periods, Patterns of Exposure, Dose Thresholds."
3. Slikker et al., "Mode of Action: Disruption of Brain Cell Replication, Second Messenger, and Neurotransmitter Systems during Development Leading to Cognitive Dysfunction — Developmental Neurotoxicity of Nicotine."
4. Slotkin, "Nicotine and the Adolescent Brain: Insights from an Animal Model."
5. Gould and Leach, "Cellular, Molecular, and Genetic Substrates Underlying the Impact of Nicotine on Learning."
6. U.S. Department of Health and Human Services, "The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General."
7. England, L.J., et al., Nicotine and the Developing Human: A Neglected Element in the Electronic Cigarette Debate. *Am J Prev Med*, 2015.
8. De Genna, N.M., et al., Prenatal tobacco exposure, maternal postnatal nicotine dependence and adolescent risk for nicotine dependence: Birth cohort study. *Neurotoxicol Teratol*, 2017.
9. Benowitz et al., "Prolonged Absorption with Development of Tolerance to Toxic Effects after Cutaneous Exposure to Nicotine."
10. Okamoto et al., "Tolerance to the Convulsions Induced by Daily Nicotine Treatment in Rats."
11. Marynak KL, Gammon DG, Rogers T, Coats EM, Singh T, King BA, "Sales of Nicotine-Containing Electronic Cigarette Products: United States, 2015", *American Journal of Public Health* 107, no. 5 (May 1, 2017): pp. 702-705. DOI: 10.2105/AJPH.2017.303660
12. Chen, J. and W.J. Millar, *Age of smoking initiation: implications for quitting*. *Health Rep*, 1998. 9(4): p. 39-46(Eng); 39-48(Fre).
13. Adelman, W.P., *Tobacco use cessation for adolescents*. *Adolesc Med Clin*, 2006. 17(3): p. 697- 717; abstract xii.
14. DiFranza, J.R. and R.J. Wellman, *A sensitization-homeostasis model of nicotine craving, withdrawal, and tolerance: integrating the clinical and basic science literature*. *Nicotine Tob Res*, 2005. 7(1): p. 9-26.
15. DiFranza, J.R. and R.J. Wellman, *Sensitization to nicotine: how the animal literature might inform future human research*. *Nicotine Tob Res*, 2007. 9(1): p. 9-20.
16. Ginzel, K.H., et al., *Critical review: nicotine for the fetus, the infant and the adolescent?* *J Health Psychol*, 2007. 12(2): p. 215-24.
17. Nelson, D.E., et al., *Long-term trends in adolescent and young adult smoking in the United States: metapatterns and implications*. *Am J Public Health*, 2008. 98(5): p. 905-15.
18. Conner, M et al., *Do electronic cigarettes increase cigarette smoking in UK adolescents? Evidence from a 12-month prospective study*, *BMJ*, 2017; 0: p. 1-9
19. Hammond, D et al., *Electronic cigarette use and smoking initiation among youth: a longitudinal cohort study*, *CMAJ*, 2017; 189: p. E1328-36